



NEWS RELEASE

# KEYTRUDA® (pembrolizumab) Approved as First Anti-PD-1 Therapy in Japan for First-Line Treatment of Patients with Advanced Non-Small Cell Lung Cancer (NSCLC) Whose Tumors Express High Levels of PD-L1

12/19/2016

KEYTRUDA Also Approved for Second-Line Treatment of Patients with Advanced NSCLC Whose Tumors Express PD-L1

Merck (NYSE:MRK), known as MSD outside the United States and Canada, today announced that KEYTRUDA® (pembrolizumab), the company's anti-PD-1 therapy, has been approved in Japan for the treatment of certain patients with PD-L1-positive unresectable advanced/recurrent non-small cell lung cancer (NSCLC) in the first- and second-line treatment settings at a fixed dose of 200 mg every three weeks. MSD will manufacture and market KEYTRUDA in Japan and will promote it with Taiho Pharmaceutical Co., Ltd.

"This approval in both the first- and second-line settings for advanced non-small cell lung cancer in patients whose tumors express PD-L1 marks an important milestone in Japan," said Dr. Roger M. Perlmutter, president, Merck Research Laboratories. "Through advanced research, Merck is demonstrating its commitment to bringing meaningful therapeutic advances to people with cancer throughout the world."

The Japanese Ministry of Health, Labour, and Welfare (MHLW) approval is supported by findings from both the KEYNOTE-024 and KEYNOTE-010 studies. KEYNOTE-024 is a randomized, open-label, phase 3 study evaluating KEYTRUDA monotherapy at a fixed dose of 200 mg compared to standard of care platinum-containing chemotherapy for the treatment of patients with both squamous and non-squamous metastatic NSCLC. The study enrolled patients who had not received prior systemic chemotherapy treatment for their metastatic disease and

whose tumors had high PD-L1 expression (as defined by a tumor proportion score [TPS] of 50 percent or more) with no EGFR or ALK aberrations. KEYNOTE-010 is an open-label, randomized, phase 2/3 trial assessing two doses of KEYTRUDA (pembrolizumab) (2 mg/kg or 10 mg/kg every three weeks) compared to docetaxel (75 mg/m<sup>2</sup> every three weeks), a standard of care chemotherapy. This study enrolled patients with any level of PD-L1 expression (as defined by a TPS of one percent or more) who had progressed following platinum-containing chemotherapy and, if appropriate, targeted therapy for EGFR or ALK genomic tumor aberrations.

The PD-L1 IHC 22C3 PharmDx™ kit made by Dako North America, Inc., an Agilent Technologies Company, was approved in Japan on Nov. 25 for use in detecting PD-L1, an immune-related biomarker expressed on some tumor cells. The diagnostic is intended to aid in identifying appropriate patients for treatment with KEYTRUDA, including previously treated patients whose tumors have any level of PD-L1 expression and previously untreated patients whose tumors have high levels of PD-L1 expression. Tumors with a TPS of less than one percent are considered to have no PD-L1 expression.

## About KEYTRUDA® (pembrolizumab)

KEYTRUDA is a humanized monoclonal antibody that works by increasing the ability of the body's immune system to help detect and fight tumor cells. KEYTRUDA blocks the interaction between PD-1 and its ligands, PD-L1 and PD-L2, thereby activating T lymphocytes which may affect both tumor cells and healthy cells.

KEYTRUDA is administered as an intravenous infusion over 30 minutes every three weeks for the approved indications. KEYTRUDA for injection is supplied in a 100 mg single use vial.

## KEYTRUDA Indications and Dosing in the U.S.

### Melanoma

KEYTRUDA is indicated for the treatment of patients with unresectable or metastatic melanoma at a dose of 2 mg/kg every three weeks until disease progression or unacceptable toxicity.

### Lung Cancer

KEYTRUDA is indicated for the first-line treatment of patients with metastatic non-small cell lung cancer (NSCLC) whose tumors have high PD-L1 expression [tumor proportion score (TPS) ≥50%] as determined by an FDA-approved test, with no EGFR or ALK genomic tumor aberrations.

KEYTRUDA (pembrolizumab) is also indicated for the treatment of patients with metastatic NSCLC whose tumors

express PD-L1 (TPS  $\geq$ 1%) as determined by an FDA-approved test, with disease progression on or after platinum-containing chemotherapy. Patients with EGFR or ALK genomic tumor aberrations should have disease progression on FDA-approved therapy for these aberrations prior to receiving KEYTRUDA.

In metastatic NSCLC, KEYTRUDA is administered at a fixed dose of 200 mg every three weeks until disease progression, unacceptable toxicity, or up to 24 months in patients without disease progression.

## Head and Neck Cancer

KEYTRUDA is indicated for the treatment of patients with recurrent or metastatic head and neck squamous cell carcinoma (HNSCC) with disease progression on or after platinum-containing chemotherapy. This indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials. In HNSCC, KEYTRUDA is administered at a fixed dose of 200 mg every three weeks until disease progression, unacceptable toxicity, or up to 24 months in patients without disease progression.

## Selected Important Safety Information for KEYTRUDA® (pembrolizumab)

KEYTRUDA can cause immune-mediated pneumonitis, including fatal cases. Pneumonitis occurred in 94 (3.4%) of 2799 patients receiving KEYTRUDA, including Grade 1 (0.8%), 2 (1.3%), 3 (0.9%), 4 (0.3%), and 5 (0.1%) pneumonitis, and occurred more frequently in patients with a history of prior thoracic radiation (6.9%) compared to those without (2.9%). Monitor patients for signs and symptoms of pneumonitis. Evaluate suspected pneumonitis with radiographic imaging. Administer corticosteroids for Grade 2 or greater pneumonitis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 or recurrent Grade 2 pneumonitis.

KEYTRUDA can cause immune-mediated colitis. Colitis occurred in 48 (1.7%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.4%), 3 (1.1%), and 4 (<0.1%) colitis. Monitor patients for signs and symptoms of colitis. Administer corticosteroids for Grade 2 or greater colitis. Withhold KEYTRUDA for Grade 2 or 3; permanently discontinue KEYTRUDA for Grade 4 colitis.

KEYTRUDA (pembrolizumab) can cause immune-mediated hepatitis. Hepatitis occurred in 19 (0.7%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.1%), 3 (0.4%), and 4 (<0.1%) hepatitis. Monitor patients for changes in liver function. Administer corticosteroids for Grade 2 or greater hepatitis and, based on severity of liver enzyme elevations, withhold or discontinue KEYTRUDA.

KEYTRUDA can cause hypophysitis. Hypophysitis occurred in 17 (0.6%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.2%), 3 (0.3%), and 4 (<0.1%) hypophysitis. Monitor patients for signs and symptoms of

hypophysitis (including hypopituitarism and adrenal insufficiency). Administer corticosteroids and hormone replacement as clinically indicated. Withhold KEYTRUDA for Grade 2; withhold or discontinue for Grade 3 or 4 hypophysitis.

KEYTRUDA can cause thyroid disorders, including hyperthyroidism, hypothyroidism, and thyroiditis. Hyperthyroidism occurred in 96 (3.4%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.8%) and 3 (0.1%) hyperthyroidism. Hypothyroidism occurred in 237 (8.5%) of 2799 patients receiving KEYTRUDA, including Grade 2 (6.2%) and 3 (0.1%) hypothyroidism. Thyroiditis occurred in 16 (0.6%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.3%) thyroiditis. Monitor patients for changes in thyroid function (at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation) and for clinical signs and symptoms of thyroid disorders. Administer replacement hormones for hypothyroidism and manage hyperthyroidism with thionamides and beta-blockers as appropriate. Withhold or discontinue KEYTRUDA for Grade 3 or 4 hyperthyroidism.

KEYTRUDA can cause type 1 diabetes mellitus, including diabetic ketoacidosis, which have been reported in 6 (0.2%) of 2799 patients. Monitor patients for hyperglycemia or other signs and symptoms of diabetes. Administer insulin for type 1 diabetes, and withhold KEYTRUDA and administer antihyperglycemics in patients with severe hyperglycemia.

KEYTRUDA can cause immune-mediated nephritis. Nephritis occurred in 9 (0.3%) of 2799 patients receiving KEYTRUDA, including Grade 2 (0.1%), 3 (0.1%), and 4 (<0.1%) nephritis. Monitor patients for changes in renal function. Administer corticosteroids for Grade 2 or greater nephritis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 nephritis

KEYTRUDA can cause other clinically important immune-mediated adverse reactions. For suspected immune-mediated adverse reactions, ensure adequate evaluation to confirm etiology or exclude other causes. Based on the severity of the adverse reaction, withhold KEYTRUDA and administer corticosteroids. Upon improvement to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Based on limited data from clinical studies in patients whose immune-related adverse reactions could not be controlled with corticosteroid use, administration of other systemic immunosuppressants can be considered. Resume KEYTRUDA (pembrolizumab) when the adverse reaction remains at Grade 1 or less following corticosteroid taper. Permanently discontinue KEYTRUDA for any Grade 3 immune-mediated adverse reaction that recurs and for any life-threatening immune-mediated adverse reaction.

The following clinically significant immune-mediated adverse reactions occurred in less than 1% (unless otherwise indicated) of 2799 patients: arthritis (1.5%), exfoliative dermatitis, bullous pemphigoid, rash (1.4%), uveitis, myositis, Guillain-Barré syndrome, myasthenia gravis, vasculitis, pancreatitis, hemolytic anemia, and partial seizures arising

in a patient with inflammatory foci in brain parenchyma.

KEYTRUDA can cause severe or life-threatening infusion-related reactions, which have been reported in 6 (0.2%) of 2799 patients. Monitor patients for signs and symptoms of infusion-related reactions, including rigors, chills, wheezing, pruritus, flushing, rash, hypotension, hypoxemia, and fever. For Grade 3 or 4 reactions, stop infusion and permanently discontinue KEYTRUDA.

Based on its mechanism of action, KEYTRUDA can cause fetal harm when administered to a pregnant woman. If used during pregnancy, or if the patient becomes pregnant during treatment, apprise the patient of the potential hazard to a fetus. Advise females of reproductive potential to use highly effective contraception during treatment and for 4 months after the last dose of KEYTRUDA.

KEYTRUDA was discontinued due to adverse reactions in 8% of 682 patients with metastatic NSCLC. The most common adverse event resulting in permanent discontinuation of KEYTRUDA was pneumonitis (1.8%). Adverse reactions leading to interruption of KEYTRUDA occurred in 23% of patients; the most common ( $\geq 1\%$ ) were diarrhea (1%), fatigue (1.3%), pneumonia (1%), liver enzyme elevation (1.2%), decreased appetite (1.3%), and pneumonitis (1%). The most common adverse reactions (occurring in at least 20% of patients and at a higher incidence than with docetaxel) were decreased appetite (25% vs 23%), dyspnea (23% vs 20%), and nausea (20% vs 18%).

It is not known whether KEYTRUDA is excreted in human milk. Because many drugs are excreted in human milk, instruct women to discontinue nursing during treatment with KEYTRUDA and for 4 months after the final dose.

Safety and effectiveness of KEYTRUDA have not been established in pediatric patients.

## Our Focus on Cancer

Our goal is to translate breakthrough science into innovative oncology medicines to help people with cancer worldwide. At Merck, helping people fight cancer is our passion and supporting accessibility to our cancer medicines is our commitment. Our focus is on pursuing research in immuno-oncology and we are accelerating every step in the journey – from lab to clinic – to potentially bring new hope to people with cancer.

As part of our focus on cancer, Merck is committed to exploring the potential of immuno-oncology with one of the fastest-growing development programs in the industry. We are currently executing an expansive research program that includes nearly 400 clinical trials evaluating our anti-PD-1 therapy across more than 30 tumor types. We also continue to strengthen our immuno-oncology portfolio through strategic acquisitions and are prioritizing the development of several promising immunotherapeutic candidates with the potential to improve the treatment of advanced cancers.

For more information about our oncology clinical trials, visit [www.merck.com/clinicaltrials](http://www.merck.com/clinicaltrials).

## About Merck

For 125 years, Merck has been a global health care leader working to help the world be well. Merck is known as MSD outside the United States and Canada. Through our prescription medicines, vaccines, biologic therapies, and animal health products, we work with customers and operate in more than 140 countries to deliver innovative health solutions. We also demonstrate our commitment to increasing access to health care through far-reaching policies, programs and partnerships. For more information, visit [www.merck.com](http://www.merck.com) and connect with us on **Twitter**, **Facebook**, **YouTube** and **LinkedIn**.

## Forward-Looking Statement of Merck & Co., Inc., Kenilworth, N.J., USA

This news release of Merck & Co., Inc., Kenilworth, N.J., USA (the “company”) includes “forward-looking statements” within the meaning of the safe harbor provisions of the U.S. Private Securities Litigation Reform Act of 1995. These statements are based upon the current beliefs and expectations of the company’s management and are subject to significant risks and uncertainties. There can be no guarantees with respect to pipeline products that the products will receive the necessary regulatory approvals or that they will prove to be commercially successful. If underlying assumptions prove inaccurate or risks or uncertainties materialize, actual results may differ materially from those set forth in the forward-looking statements.

Risks and uncertainties include but are not limited to, general industry conditions and competition; general economic factors, including interest rate and currency exchange rate fluctuations; the impact of pharmaceutical industry regulation and health care legislation in the United States and internationally; global trends toward health care cost containment; technological advances, new products and patents attained by competitors; challenges inherent in new product development, including obtaining regulatory approval; the company’s ability to accurately predict future market conditions; manufacturing difficulties or delays; financial instability of international economies and sovereign risk; dependence on the effectiveness of the company’s patents and other protections for innovative products; and the exposure to litigation, including patent litigation, and/or regulatory actions.

The company undertakes no obligation to publicly update any forward-looking statement, whether as a result of new information, future events or otherwise. Additional factors that could cause results to differ materially from those described in the forward-looking statements can be found in the company’s 2015 Annual Report on Form 10-K and the company’s other filings with the Securities and Exchange Commission (SEC) available at the SEC’s Internet site ([www.sec.gov](http://www.sec.gov)).

Please see Prescribing Information for KEYTRUDA (pembrolizumab) at [http://www.merck.com/product/usa/pi\\_circulars/k/keytruda/keytruda\\_pi.pdf](http://www.merck.com/product/usa/pi_circulars/k/keytruda/keytruda_pi.pdf) and

Patient Information/Medication Guide for KEYTRUDA at [http://www.merck.com/product/usa/pi\\_circulars/k/keytruda/keytruda\\_mg.pdf](http://www.merck.com/product/usa/pi_circulars/k/keytruda/keytruda_mg.pdf).

Merck

Media:

Pamela Eisele, 267-305-3558

Courtney Ronaldo, 908-236-1108

or

Investors:

Teri Loxam, 908-740-1986

Amy Klug, 908-740-1898