



NEWS RELEASE

New Studies Investigating the Use of KEYTRUDA® (pembrolizumab), Merck's Anti-PD-1 Therapy, in Advanced Melanoma Compared to Chemotherapy, in Classical Hodgkin Lymphoma and in Triple Negative Breast Cancer, to be Presented for the First Time

11/6/2014

By Year End, Data on KEYTRUDA in Seven Tumor Types Will Have Been Presented

WHITEHOUSE STATION, N.J.--(BUSINESS WIRE)--Merck (NYSE:MRK), known as MSD outside the United States and Canada, announced today that in November and December, data will be presented for the first time investigating the use of KEYTRUDA® (pembrolizumab) – the company's anti-PD-1 therapy – in advanced melanoma in comparison to chemotherapy, and in relapsed/refractory classical Hodgkin Lymphoma (cHL) as well as advanced triple negative breast cancer (TNBC). These studies will be presented in oral sessions at the Society for Melanoma Research (SMR) 2014 International Congress, Nov. 13 – 16, the 56th American Society of Hematology (ASH) Annual Meeting, Dec. 6 – 9, and the San Antonio Breast Cancer Symposium (SABCS), Dec. 9 – 13, respectively. By the end of 2014, data on anti-tumor activity of KEYTRUDA as monotherapy will have been presented in seven different tumor types.

“Our focus on evaluating KEYTRUDA as a monotherapy is enabling Merck to rapidly advance the clinical development of our anti-PD-1 therapy,” said Dr. Roy Baynes, senior vice president, Global Clinical Development, Merck Research Laboratories. “We look forward to the first comparative data in advanced melanoma, and new data in advanced triple negative breast cancer and classical Hodgkin Lymphoma being presented. We continue to expand our immuno-oncology clinical program with KEYTRUDA both as monotherapy and in combination with

other agents.”

KEYTRUDA is indicated in the United States at a dose of 2 mg/kg every three weeks for the treatment of patients with unresectable or metastatic melanoma and disease progression following ipilimumab and, if BRAF V600 mutation positive, a BRAF inhibitor. This indication is approved under accelerated approval based on tumor response rate and durability of response. An improvement in survival or disease-related symptoms has not yet been established. Continued approval for this indication may be contingent upon verification and description of clinical benefit in the confirmatory trials.

Merck data to be presented at SMR 2014 include:

- Late Breaker Oral Presentation: A Randomized Controlled Comparison of Pembrolizumab (MK-3475) and Chemotherapy in Patients with Ipilimumab-Refractory Melanoma (KEYNOTE-002). A. Ribas. Sunday, November 16, 8:50 AM-10:25 AM CET. Location: Kongresshaus, First Floor.

Merck data to be presented at the ASH Annual Meeting include:

- (Abstract #290) Oral Presentation: PD-1 Blockade with the Monoclonal Antibody Pembrolizumab (MK-3475) in Patients with Classical Hodgkin Lymphoma After Brentuximab Vedotin Failure: Preliminary Results From a Phase 1b Study (KEYNOTE-013). C. Moskowitz. Monday, December 8, 7:15 AM-7:30 AM PST. Location: Moscone Center, West Building, 3009-3011-3022-3024.

Merck data to be presented at SABCS include:

- (Abstract #S1-09) Oral Presentation: A Phase 1b Study (KEYNOTE-012) of Pembrolizumab (MK-3475) in Patients with Advanced Triple-Negative Breast Cancer. R. Nanda. Wednesday, December 10, 10:45 AM-11:00 AM CST. Location: Henry B. Gonzalez Convention Center, Hall D.

For more information including a complete list of abstract titles, please visit the SMR 2014 website at www.melanomacongress.com, the ASH Annual Meeting website at www.hematology.org/Annual-Meeting, and the SABCS website at www.sabcs.org.

About KEYTRUDA (pembrolizumab)

KEYTRUDA (pembrolizumab) is a humanized monoclonal antibody that blocks the interaction between PD-1 and its ligands, PD-L1 and PD-L2. By binding to the PD-1 receptor and blocking the interaction with the receptor ligands, KEYTRUDA releases the PD-1 pathway-mediated inhibition of the immune response, including the anti-tumor immune response.

Selected Important Safety Information for KEYTRUDA

Pneumonitis occurred in 12 (2.9%) of 411 patients with advanced melanoma receiving KEYTRUDA (the approved indication in the United States), including Grade 2 or 3 cases in 8 (1.9%) and 1 (0.2%) patients, respectively. Monitor patients for signs and symptoms of pneumonitis. Evaluate suspected pneumonitis with radiographic imaging. Administer corticosteroids for Grade 2 or greater pneumonitis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 pneumonitis.

Colitis (including microscopic colitis) occurred in 4 (1%) of 411 patients, including Grade 2 or 3 cases in 1 (0.2%) and 2 (0.5%) patients respectively, receiving KEYTRUDA. Monitor patients for signs and symptoms of colitis. Administer corticosteroids for Grade 2 or greater colitis. Withhold KEYTRUDA for Grade 2 or 3; permanently discontinue KEYTRUDA for Grade 4 colitis.

Hepatitis (including autoimmune hepatitis) occurred in 2 (0.5%) of 411 patients, including a Grade 4 case in 1 (0.2%) patient, receiving KEYTRUDA. Monitor patients for changes in liver function. Administer corticosteroids for Grade 2 or greater hepatitis and, based on severity of liver enzyme elevations, withhold or discontinue KEYTRUDA.

Hypophysitis occurred in 2 (0.5%) of 411 patients, including a Grade 2 case in 1 and a Grade 4 case in 1 (0.2% each) patient, receiving KEYTRUDA. Monitor for signs and symptoms of hypophysitis. Administer corticosteroids for Grade 2 or greater hypophysitis. Withhold KEYTRUDA for Grade 2; withhold or discontinue for Grade 3; and permanently discontinue KEYTRUDA for Grade 4 hypophysitis.

Nephritis occurred in 3 (0.7%) patients receiving KEYTRUDA, consisting of one case of Grade 2 autoimmune nephritis (0.2%) and two cases of interstitial nephritis with renal failure (0.5%), one Grade 3 and one Grade 4. Monitor patients for changes in renal function. Administer corticosteroids for Grade 2 or greater nephritis. Withhold KEYTRUDA for Grade 2; permanently discontinue KEYTRUDA for Grade 3 or 4 nephritis.

Hyperthyroidism occurred in 5 (1.2%) of 411 patients, including Grade 2 or 3 cases in 2 (0.5%) and 1 (0.2%) patients respectively, receiving KEYTRUDA. Hypothyroidism occurred in 34 (8.3%) of 411 patients, including a Grade 3 case in 1 (0.2%) patient, receiving KEYTRUDA. Thyroid disorders can occur at any time during treatment. Monitor patients for changes in thyroid function (at the start of treatment, periodically during treatment, and as indicated based on clinical evaluation) and for clinical signs and symptoms of thyroid disorders. Administer corticosteroids for Grade 3 or greater hyperthyroidism. Withhold KEYTRUDA for Grade 3; permanently discontinue KEYTRUDA for Grade 4 hyperthyroidism. Isolated hypothyroidism may be managed with replacement therapy without treatment interruption and without corticosteroids.

Other clinically important immune-mediated adverse reactions can occur. The following clinically significant,

immune-mediated adverse reactions occurred in less than 1% of patients treated with KEYTRUDA: exfoliative dermatitis, uveitis, arthritis, myositis, pancreatitis, hemolytic anemia, partial seizures arising in a patient with inflammatory foci in brain parenchyma, adrenal insufficiency, myasthenic syndrome, optic neuritis, and rhabdomyolysis.

For suspected immune-mediated adverse reactions, ensure adequate evaluation to confirm etiology or exclude other causes. Based on the severity of the adverse reaction, withhold KEYTRUDA and administer corticosteroids. Upon improvement of the adverse reaction to Grade 1 or less, initiate corticosteroid taper and continue to taper over at least 1 month. Restart KEYTRUDA if the adverse reaction remains at Grade 1 or less. Permanently discontinue KEYTRUDA for any severe or Grade 3 immune-mediated adverse reaction that recurs and for any life-threatening immune-mediated adverse reaction.

Based on its mechanism of action, KEYTRUDA may cause fetal harm when administered to a pregnant woman. If used during pregnancy, or if the patient becomes pregnant during treatment, apprise the patient of the potential hazard to a fetus. Advise females of reproductive potential to use highly effective contraception during treatment and for 4 months after the last dose of KEYTRUDA.

For the treatment of advanced melanoma, KEYTRUDA was discontinued for adverse reactions in 6% of 89 patients who received the recommended dose of 2 mg/kg and 9% of 411 patients across all doses studied. Serious adverse reactions occurred in 36% of patients receiving KEYTRUDA. The most frequent serious adverse drug reactions reported in 2% or more of patients were renal failure, dyspnea, pneumonia, and cellulitis.

The most common adverse reactions (reported in $\geq 20\%$ of patients) were fatigue (47%), cough (30%), nausea (30%), pruritus (30%), rash (29%), decreased appetite (26%), constipation (21%), arthralgia (20%), and diarrhea (20%).

The recommended dose of KEYTRUDA is 2 mg/kg administered as an intravenous infusion over 30 minutes every three weeks until disease progression or unacceptable toxicity. No formal pharmacokinetic drug interaction studies have been conducted with KEYTRUDA. It is not known whether KEYTRUDA is excreted in human milk. Because many drugs are excreted in human milk, instruct women to discontinue nursing during treatment with KEYTRUDA. Safety and effectiveness of KEYTRUDA have not been established in pediatric patients.

Our Focus on Cancer

Our goal is to translate breakthrough science into biomedical innovations to help people with cancer worldwide. For Merck Oncology, helping people fight cancer is our passion, supporting accessibility to our cancer medicines is our commitment, and pursuing research in immuno-oncology is our focus to potentially bring new hope to people with cancer. For more information about our oncology clinical trials, visit www.merck.com/clinicaltrials.

About Merck

Today's Merck is a global healthcare leader working to help the world be well. Merck is known as MSD outside the United States and Canada. Through our prescription medicines, vaccines, biologic therapies and animal health products, we work with customers and operate in more than 140 countries to deliver innovative health solutions. We also demonstrate our commitment to increasing access to healthcare through far-reaching policies, programs and partnerships. For more information, visit www.merck.com and connect with us on **Twitter**, **Facebook** and **YouTube**.

Forward-Looking Statement

This news release includes "forward-looking statements" within the meaning of the safe harbor provisions of the United States Private Securities Litigation Reform Act of 1995. These statements are based upon the current beliefs and expectations of Merck's management and are subject to significant risks and uncertainties. There can be no guarantees with respect to pipeline products that the products will receive the necessary regulatory approvals or that they will prove to be commercially successful. If underlying assumptions prove inaccurate or risks or uncertainties materialize, actual results may differ materially from those set forth in the forward-looking statements.

Risks and uncertainties include, but are not limited to, general industry conditions and competition; general economic factors, including interest rate and currency exchange rate fluctuations; the impact of pharmaceutical industry regulation and healthcare legislation in the United States and internationally; global trends toward healthcare cost containment; technological advances, new products and patents attained by competitors; challenges inherent in new product development, including obtaining regulatory approval; Merck's ability to accurately predict future market conditions; manufacturing difficulties or delays; financial instability of international economies and sovereign risk; dependence on the effectiveness of Merck's patents and other protections for innovative products; and the exposure to litigation, including patent litigation, and/or regulatory actions.

Merck undertakes no obligation to publicly update any forward-looking statement, whether as a result of new information, future events or otherwise. Additional factors that could cause results to differ materially from those described in the forward-looking statements can be found in Merck's 2013 Annual Report on Form 10-K and the company's other filings with the Securities and Exchange Commission (SEC) available at the SEC's Internet site (www.sec.gov).

#

Please see Prescribing Information for KEYTRUDA (pembrolizumab) at http://www.merck.com/product/usa/pi_circulars/k/keytruda/keytruda_pi.pdf and the Medication Guide for KEYTRUDA at http://www.merck.com/product/usa/pi_circulars/k/keytruda/keytruda_mg.pdf.

KEYTRUDA® is a registered trademark of Merck & Co., Inc., Whitehouse Station, N.J., USA

Merck

Media:

Pamela Eisele, 267-305-3558

Claire Mulhearn, 908-236-1118

or

Investor:

Joseph Romanelli, 908-423-5185

Justin Holko, 908-423-5088